PART I. SUBJECT INDEX

Abandoned mines, in Ohio, Iowa and West Virginia, discharging acid into streams, now sealed, Denmead, Talbott, 31-32.
Abnormalities of fish in polluted waters (disc.), 203.
Abscesses in new hatchery disease of trout, pathology, symptomatology, etiology and epidemiology discussed, Belding, David L., and Merrill, Beulah, 76-84
Abundance of fish erroneously figured: see **Statistics.**
of suckers indicative of reduction in numbers of sport fish, Huntsman, A. G., 152-156.
Acantholeberis curvirostris as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Acetic acid: (See also **Glacial acetic acid.**)
treatment, experimental work, Wolf, Louis E., 88-100.
Advisory Committee of the Secretary of Commerce, recommendations, Bell, Frank T., 247-248.
Air-dried salmon meal found superior to commercial salmon meal as food for young salmon, indicating that composition and method of manufacture are important, Donaldson, Lauren R., 169.
Airplane method of stocking speckled trout, Prevost, Gustave, 277-278.
Alabama, adoption of commission form of conservation administration; what was Department of Game and Fisheries is now Department of Conservation of Game, Fish and Sea Foods, with powers to fix open seasons and bag limits on all game and creel limits on fish; bill passed prohibiting sale of bass, Quinn, I. T. (disc.), 42.
Alewives as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 381-382.
Algae affected by copper sulphate, about two parts per million, but rapidly recovered, Hazzard, A. S. (disc.), 114.
as affected by concentrations of copper sulphate in Lake Jesse, Smith, M. W., 106.
malodorous, control in Madison lakes, Domogalla, Bernhard, 115-120.
Alonella nana, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Ambloplites rupestris, analysis of catch in Fife Lake, Eschmeyer, R. W., 210-211.
Ameiurus melas specimens taken at four Buckeye Lake stations, stomach contents tabulated, 63.
nebulosus, A. natalis, analysis of catch in Fife Lake, Eschmeyer, R. W., 212.
nebulosus surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
American Fish Policy Committee, report, Wickliff, E. L., 27-29.
American Fisheries Society, A research Committee recommended to see what should most urgently be done and assign it to the proper agency (either State or Federal), Wiebe, A. H., 37.
certificate of incorporation. 405.
committees 1935-1936, list of members, 4-5.
constitution and by-laws, 406-409.
list of members, honorary: 410; patrons; 410-411; life; 412; active: 413-427; state membership: 427-428; clubs, dealers, etc.: 428-429; libraries: 429; corresponding members: 429.
officers 1934-1935; 1935-1936, 3.
American Wildlife Institute, resolve by American Fisheries Society to pledge its hearty cooperation, 444.
sportsmen organized by its recent formation, Denmead, Talbott, 41.
Amphipods, Hyalella knickerbockeri: see **Hyalella.**

Anabena control in Madison lakes, Domogalla, Bernhard, 115-120.
Angling, report Division of, Denmead, Talbott, 40-42.
statistics from Furnace Brook "test stream" suggested a later season and smaller creel limit for trout fishing in northern states, Lord, Russell F., 224-231.
Anguilla rostrata surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Anuraea cochlearis, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Aphanizomenon control in Madison lakes, Domogalla, Bernhard, 115-120.
controlled with copper sulphate, about five parts per million, without any deleterious effect on invertebrate or fish life of the lake, Aitken, W. W. (disc.), 113.
Aphanothece (clathrata?) as affected by concentrations of copper sulphate in Lake Jesse, Smith, M. W., 106.
Aplites salmoides, analysis of catch in Fife Lake, Eschmeyer, R. W., 212.
specimens taken at eight Buckeye Lake stations, stomach contents tabulated, 66.
Appendix, 403.
Aquarium observations on predatory fish add red-bellied dace and blunt-nosed minnow to list of forage fish, Cooper, Gerald P., 142.
Aquatic biology and physics, report Vice-president Division of, Wiebe, A. H., 36-37.
insects surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
plants, distribution and abundance in Fish Lake, tabulated, Hazzard, A. S., 124.
Aquicultural suggestions, Langlois, T. H., 284-287.
Arkansas State Fish Commission reports development of a very efficient method for solving the problem of increasing costs of fish foods, method described, Fiedler, R. H., and Samson, V. J., 390.
Artificial fly bait vs. other bait, analysis of catch in Fife Lake, Eschmeyer, R. W., 215-217.
propagation and stocking in the national forests, Davis, H. S., 236-237.
Asterionella formosa, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Atlantic Ocean, transgression of marginal waters over the Scotian Shelf, Leim, A. H., and Hachey, H. B., 279-283.
salmon parr, growth habits, Belding, David L., 157-160.
Auditing Committee report, Denmead, Talbott, 43.
Bacteria, Gram-positive diplo-bacilli, present in new hatchery disease of trout, Belding, David L., and Merrill, Beulah, 76-84.
Bacterial gill disease with lesions comparable to those described by Davis, but with distinctly different causative organism, described, Fish, Frederic F., 85-87, 4 plates.
Bait, types used during census of Fife Lake, Eschmeyer, R. W., 214-217.
Baits, artificial more in use than formerly, Denmead, Talbott, 41.
Bass, black, protective legislation, Amsler, Guy, 38-39.
fry in aquaria, eaten by forage fish, including red-bellied dace and blunt-nosed minnow, Cooper, Gerald P., 142.
largemouth (see also **Aplites** salmoides), food habits in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 59-60; tabulated, 66.
smallmouth (see also **Micropterus** dolomieu), propagated in troughs at South Otselic, Kingsbury, O. R., and Royce, Wm. F., 309-312.
striped, sale in California prohibited by law, Amsler, Guy, 39.
white, closed season abolished in Michigan, Amsler, Guy, 38.
white, diet in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 60; tabulated, 68.
Beaver-trout management program in Michigan, Bradt, Glenn W., 253-257.

Beef liver and air-dried salmon meal produced very rapid growth of young king salmon, with lowest mortality rate and minimum cost in experimental feeding, Donaldson, Lauren R., 165-170.
meat, hearts, kidneys, livers, lungs, melts (spleen), analysis of protein, fat, carbohydrate and ash content tabulated, 397.
Bell, Frank T., elected President American Fisheries Society, 45-46.
Bicarbonate compounds important factors in causing obnoxious water growths in Madison lakes, Domogalla, Bernhard, 115.
Black bullhead, diet, Ewers, Lela A., and Boesel, M. W., 58; tabulated, 63.
crappie: see **Crappie,** black.
Blebs: see **Lesions.**
Blood meal (dried), analysis of protein, fat, carbohydrates and ash content tabulated, 397.
Bluegill (see also **Helioperca** macrochira), diet, Ewers, Lela A., and Boesel, M. W., 59; tabulated, 65.
Blunt-nosed minnow, diet, Ewers, Lela A., and Boesel, M. W., 57; tabulated, 61.
life history observations bearing on their culture, Cooper, Gerald P., 139-140.
relative growth rate of male and female, Cooper, Gerald P., 137.
Bosmina longirostris, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Bottom of ponds, straight sand and gravel do not roil; addition of clay in conjunction with planting of crayfish for control of vegetation recommended, Langlois, T. H., 193.
Brazil, northeastern, fisheries investigations , Ihering, Rodolpho von, and Wright, Stillman, 267-271.
Breeding habits of the stone roller minnow, Campostoma anomalum Rafinesque, Smith, Osgood R., 148-151.
British Columbia pond culture objectives, Mottley, C. McC., 172-178.
Brook silverside, diet, Ewers, Lela A., and Boesel, M. W., 59; tabulated, 64.
Buckeye Lake, diet of various fish species, Ewers, Lela A., and Boesel, M. W., 57-69.
Bullhead: see **Ameiurus** spp. and **Black bullhead.**
Burbot in Minnesota lakes (disc.), 274.
Bureau of Fisheries reports increased production and distribution of fish and eggs, including fry hatched on a cooperative basis in State hatcheries, approximated at 5,000,000,000 in comparison with an output of 3,258,000,000 for the previous year, Jackson, Charles E., in report on Committee on Relations with Federal, Provincial and State governments, 25.
By-laws, American Fisheries Society, 406-409.
CCC men: see **Civilian** Conservation Corps.
Calcium requirements of trout, Tunison, A. V., and McCay, C. M., 359-361.
California hatcheries, experiments in trout feeding, Wales, Joseph H., 305-308.
protective legislation, Amsler, Guy, 38-39.
two small mountain lakes utilized to determine the annual production of this type of water; experimental streams set aside for intensive study, including losses occurring among fish of various sizes after planting and extent to which natural food can be utilized without danger of diminishing returns; life-history of steelhead trout; effects of stream obstructions and diversions on migratory fish, Davis, H. S., 239.
Callinectes sapidus, nutritive value, Watson, Vernon K., and Fellers, Carl R., 342-349.
Cambarus rusticus Girard, habits in Ohio fish ponds, Langlois, T. H., 189-192.
Campostoma anomalum Refinesque, breeding habits, Smith, Osgood R., 148-151.

Carbon and nitrogen percentages in Potamogeton americanus, Ceratophyllum demersum and green vegetation, Meehean, O. Lloyd, 186.
Carbon dioxide, effect on development of trout eggs, Surber, Eugene W., 194-203.
Carp, food taken under natural conditions, Ewers, Lela A., and Boesel, M. W., 57; tabulated, 61.
Cary, Guy, Life Member American Fisheries Society, was reported deceased in the 1934 Transactions, error corrected, 47.
Casting vs. other methods of fishing, analysis of catch in Fife Lake, Eschmeyer, R. W., 214-217.
Catfish: see **Ameiurus** nebulosus and **Tadpole** catfish.
Catostomus commersonii, analysis of catch in Fife Lake, Eschmeyer, R. W., 212.
in relation to Salmo salar and Salvelinus fontinalis in Margaree River and tributaries, Cape Breton Island, Nova Scotia, Huntsman, A. G., 152-156.
surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Census of fishing on Fife Lake, Michigan, taken by CCC men, Eschmeyer, R. W., 207-223.
Ceratophyllum demersum, chemical analysis showed a percentage of 42.38 carbon and 1.9 nitrogen, Meehean, O. Lloyd, 186.
distribution and abundance in Fish Lake, Utah, Hazzard, A. S., 124.
Ceriodaphnia reticulata in diet of bass and sunfish, Ewers, Lela A., and Boesel, M. W., 59-60.
Chanel cat, the small series studied showed remains of midge larvae and pupae, Ewers, Lela A., and Boesel, M. W., 58.
Chara sp., distribution and abundance in Fish Lake, Hazzard, A. S., 124.
Chemical treatment of waters of Madison lakes over eleven-year period; its effect on fish and fish foods, Domogalla, Bernhard, 115-120.
Cheonda hydrophlox introduced into Fish Lake by fishermen as live bait, Hazzard, A. S., 127.
Chironomid larvae and pupae as food for trout in lakes of the Klamath River watershed, Doudoroff, Peter, 130.
surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Chronic infectious disease of hatchery trout, hitherto undescribed, pathology, symptomatology, and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
Chroococcus, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Chrosomus eos, propagation experiments, Cooper, Gerald P., 132-142.
Chub (see also **Semotilus** atromaculatus), spawning habits, Schultz, Leonard P., 143-147.
sucker, life history observations bearing on their culture, Cooper, Gerald P., 141.
unrestricted use as live bait liable to prove distinct annoyance to fly fishermen if they become numerous after being thus introduced, as in Fish Lake, Hazzard, A. S., 127.
Chydorus sphaericus, an affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
in stomachs of blunt-nosed minnows, Ewers, Lela A., and Boesel, M. W., 57.
Civilian Conservaiton Corps, fishing census taken on Fife Lake, Eschmeyer, R. W., 207-223.
progress made in stream improvement, Greeley, John R., 316-321.
Cladocera, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
in diet of various species of fish in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 57-69.
Cladophora control in Madison lakes, Domogalla, Bernhard, 115-120.

Clam products as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 385-386.
Clear Lake, California, taking of crappie at any time authorized, Amsler, Guy, 38.
Closterium, as affected by copper sulphate in Lake Jesse, Nova Scotia, Smith, M. W., 107.
Cobb, Eben W., elected Librarian American Fisheries Society, 1935, 45-46.
Cold and ice in inland waters, winter research needed to determine extent of their affect on fish life, Hubbs, Carl L., and Trautman, Milton B., 51-56.
Cold storage warehouses and freezing plants in the United States and Alaska, tabulated, 391.
Coldwater Creek, Winneshiek County, Iowa, construction faults, ineffectual devices and misapplied principles in stream improvement shown, Aitken, W. W., 322-323, 2 plates.
Commercial food, diet of sufficient amount to produce growth comparable to that obtained in other lots tested resulted in excessive mortality after ten days; when fed at the recommended low level, unsatisfactory growth of the brook trout was noted, Deuel, Charles R., 163.
Commissao Technica de Pisciculture, Fortaleza, Ceara, Northeast Brazil, fishery investigations, Ihering, Rodolpho von, and Wright, Stillman, 267-271.
Commission form of administration established in states of Alabama, Florida, Maryland, New Hampshire, Rhode Island, South Carolina and Tennessee, Amsler, Guy, 39.
Committee (see also **Foreign** relations), appointment, 42.
on Common and Scientific Names of Fishes, members elected: Hubbs, Carl L., Hildebrand, Samuel F., Harkness, Wm. J. K., Snyder, John O., Chute, Walter H., and Leim, A. H., 46.
on Time and Place, motion that the 1936 meeting be held the first week in September at Grand Rapids, Michigan carried unanimously, 44-45.
Common and scientific names of fishes, Committee: see **Committee.**
Common names of important North American fishes, report committee on, Hubbs, Carl, L., 26-27.
sunfish, diet, Ewers, Lela A., and Boesel, M. W., 59; tabulated, 65.
Constitution and by-laws, 406-409.
Cooperation in fishery investigations, Bell, Frank T., 248-252.
Connecticut, creation of Interstate Sanitation District and Commission with the states of New York and New Jersey, authorized by Resolution adopted by Congress (S. J. Res. 159), 32-33.
Conorhynchus conirostris from Sao Francisco River, Brazil, Ihering, Rodolpho von, and Wright, Stillman, 268.
Copepod infection of speckled trout, Savage, James, 334-339, 1 figure.
Copepoda as diet of various species of fish in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 57-69.
Copepods as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Copper sulphate, about five parts per million, used without deleterious effect on invertebrate life or fish life, yet controlled Aphanizomenon and Microcystis, Aitken, W. W. (disc.), 113.
about two parts per million, killed the snails, but also the trout, Hazzard, A. S. (disc.), 114.
and sodium arsenate vs. crayfish for elimination of vegetation in Ohio fish ponds, Langlois, T. H., 192.
causes death of fish apparently by a precipitation of the mucus on the gills, with subsequent suffocation, rather then by internal poisoning, 111; experiments in Lake Jesse and results obtained, Smith, M. W., 101-113.
spraying method most effective and economical in keeping obnoxious algae under control; eleven-year treatment in Madison lakes did not pre-

vent luxurious growth of rooted water weeds, Domogalla, Bernhard, 115-120.
Corethra larvae surviving addition of copper sulphate in Lake Jesse, Smith, M. W.. 109.
Cortland Diets, feeding experiments with brook trout at New York State Hatchery, Gloversville, Deuel, Charles R., 161-163.
Cosmarium, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Costia nextarix causing costiasis, identification and methods of treatment, Savage, James, 332-333.
Cottonseed meal in trout diets, Deuel, Charles R., 161-163.
Council American Fisheries Society, Report of 1935 meeting, Gordan, Seth, 21.
Crabs, Callinectes sapidus and Platyonichus ocellatus, nutritive value, Watson, Vernon K., and Fellers, Carl R., 342-349.
Crappie (see also **Pomoxis** sparoides and **White** crappie), black live water mites found in intestine; at other times the mites were digested; suggested preference of larger crappie for insect larvae, yet instance showed the largest with entire stomach and entire intestine completely filled with cereodaphnia, Wiebe, A. H. (disc.), 69-70.
open season in California changed from May 1 to November 30 and taking of them at any time authorized in District 4¼ and Clear Lake, Amsler, Guy, 38.
Crayfish may be effectively eliminated from a pond by draining in early September, before the pairing season; notes on the habits of Cambarus rusticus in Ohio fish ponds, Langlois, T. H., 189-192.
Creek club: see **Semotilus** atromaculatus.
Creel census on Lake Fife, Michigan, Eschmeyer, R. W., 207-223.
Cristovomer namaycush in Fish Lake, Utah, Hazzard, A. S., 126.
Crustacea in diet of various species of fish in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 57-69.
Crustacean fish food, hard water may not be essential to heavy yield, Hazzard, A. S., 125.
Cyclops americanus in diet of blue gill and yellow perch in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 59-60.
viridis, as affected by copper sulphate in Lake Jesse, Smith M. W., 108.
Cyprinus carpio specimens taken at two Buckeye Lake stations, stomach contents tabulated, 61.
Dairy products, analysis of protein, fat, carbohydrates and ash content tabulated, 397.
as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 386.
Dams constructed by CCC men in New York State, Greeley, John R., 317-318.
Damselfly nymphs conspicuous in diet of largemouth bass in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 60.
Daphnia pulex, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Death of fish by copper sulphate caused by a precipitation of the mucus on the gills, with subsequent suffocation, rather than by internal poisoning, Smith, M. W., 111.
Deformity of the spinal column in rainbow and brook trout caused by a concentration of 55-78.5 parts per million of carbon dioxide, Surber, Eugene W., 201; 203.
Delaware, extension of closed season on black bass, and sale of black bass in state prohibited by law, Amsler, Guy, 38-39.
Dentition of the pike, Trautman, Milton B., and Hubbs, Carl L., 261-266, 2 figures.
Desmidae, Hyalotheca, present in plankton samples in waters of Jesse Lake, protected from copper sulphate by its gelatinous sheath, Smith, M. W., 106.

Development of trout eggs as effected by carbon dioxide, Surber, Eugene W., 194-203.
Diaphanosoma, found absent in Lake Jesse treated with copper sulphate, and present in control lake, Smith, M. W., 109.
leuchtenbergianum in diet of white bass in Buckeye Lake; found to be the most abundant food organism in white crappie in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 59, 60.
Diaptomus minutus, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Diatoms as affected by concentrations of copper sulphate in Lake Jesse, Smith, M. W., 107.
Diet of salmon by-products in experimental feeding of young king salmon and results obtained, Donaldson, Lauren R., 165-170.
of various species of fish in Buckeye Lakes, Ewers, Lela A., and Boesel, M. W., 57-69.
Diets, analyses of commercial feed combinations, tabulated, 370.
experiments in feeding brook trout at New York State Hatchery at Gloversville, results tabulated, Deuel, Charles R., 163.
Digestion and utilization of fat by trout, Tunison, A. V., and McCay, C. M., 361-375.
Dinobryon, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Dipper for trout feeding, new type, Wales, Joseph H., 307.
Disease of fish, copepod infection of speckled trout, Savage, James, 334-339, 1 figure.
hatchery disease of Salmonidae, preliminary report of pathology of lesions, symptomatology, etiology and epidemiology, Belding, David L., and Merrill, Beulah, 76-84.
octomitiasis in Lake Edward, Quebec, Richardson, L. R., 290-292.
protozoan, costiastis, identification and treatment, Savage, James, 332-333.
use of potassium permanganate, experimental work, Wolf, Louis E., 88-100.
Western type of bacterial gill disease, with lesions comparable to those of the Eastern type described by Davis, but with distinctly different causative organism, is described, Fish, Frederic F., 85-87, 4 plates.
Dorosoma cepedianum (Gizzard shad), seven specimens 22-31 mm. total length taken at three Buckeye Lake stations showed a stomach contents of 14.29 per cent Cladocera and 85.71 per cent algae and debris, Ewers, Lela A., and Boesel, M. W., 61.
Dried skim milk: see **Milk,** dried skim.
Dry meals gave satisfactory results in experiments on diet of brook trout and is recommended for fish from one and one-half inches on, Deuel, Charles R., 163.
Drying of meal for fish food, steam-dried versus air-dried method, Donaldson, Lauren R., 170.
Ecology, differing environmental conditions affect the growth rate of salmon parr, Belding, David L., 157-160.
of aquatic habitats, Meehean, O. Lloyd, 184-188.
Economy vs. results obtained in feeding of fish (disc.), Tunison, A. V., 164.
Eel: see **Anguilla** rostrata.
Eggs of club layed in shallow water within two feet of shore line, Schultz, Leonard P., 146-147.
of Prochilodus extruded 6-10 hours after injection of 2 to 4 hypophyses, Ihering, Rodolpho von, and Wright, Stillman, 270.
of Tetragonopterus extruded 6-10 hours after injection of ½ hypophysis, Ihering, Rodolpho von, and Wright, Stillman, 270.
of trout, effects of carbon dioxide on development, Surber, Eugene W., 194-203.
Embryonic rainbow and brook trout as affected by carbon dioxide, Surber, Eugene W., 194-203.
Emmons, H. Nelson, deceased, 47

Entomostraca, mostly Cladocera; also Copepoda and Ostracoda, as food for trout in lakes of the Klamath River watershed, Doudoroff, Peter, 130.
Environmental conditions, effect on growth of salmon parr, Belding, David L., 157-160.
Epischura lacustris, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Erimyzon sucetta kennerlii, propagation experiments, Cooper, Gerald P., 132-142.
specimens taken at two Buckeye Lakes stations, stomach contents tabulated, 61.
Eriocaulon articulare found alive in waters with addition of copper sulphate, Smith, M. W., 108.
Esox lucius, analysis of catch in Fife Lake, Eschmeyer, R. W., 212.
lucius, E. masquinongy masquinongy, E. m. immaculatus, and E. niger, dentition, Trautman, Milton B., and Hubbs, Carl, L., 261-266, 2 figures.
Etiology of new hatchery disease of trout, Belding, David L., and Merrill, Beulah, 76-84.
Eucyclops agilis, E. prasinus, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Eupomotis gibbosus, analysis of catch in Fife Lake, Eschmeyer, R. W., 212.
specimens taken from Buckeye Lake, stomach contents tabulated, 65.
Exophthalmos in new disease of trout, Belding, David L., and Merrill, Beulah, 76-84.
Eyes gouged out during spawning, Smith, Osgood R., 151.
Fat, digestion and utilization by trout, Tunison, A. V., and McCay, C. M., 361-366.
Federal and State cooperation in fishery investigations, Bell, Frank T., 246-252.
Provincial and State Governments, Committee on Relations, report 1934-1935, Jackson, Charles E., 24-26.
Feeding experiments, technique in calculating weight of experimental groups; weight of trout per cubic foot of water; amounts to feed trout, Tunison, A. V., and McCay, C. M., 373-374.
habits of salmon parr in Margaree River and tributaries, Cape Breton Island, Nova Scotia, Huntsman, A. G., 152-156.
of brook trout, no concern need be felt in changing abruptly from a meat to a dry food diet since experiments show a resultant acceleration rather than retardation in growth, Deuel, Charles R., 163.
of pond fish at Tulsa municipal hatchery, Oklahoma, Aldrich, A. D., 179-183.
of trout of California hatcheries, Wales, Joseph H., 305-308.
practices and fish hatchery foods, survey, Fiedler, R. H., and Samson, V. J., 376-400.
Fertilizer effect on oxygen supply, Wiebe, A. H. (disc.), 36-37.
Fertilizing substances, dispersal in ponds, Meehean, O. Lloyd, 184-188.
Fesler, D. F., deceased, 47
Fiedler, R. H., elected Vice-President Division of Commercial Fishing, 1935, 45-46.
Field work in winter should be carried on to determine effect of cold and ice conditions on fish life, Hubbs, Carl L., and Trautman, M. B., 51-56.
Fife Lake creel census, Eschmeyer, R. W., 207-223.
Fish culture, report Vice-President Division of, 1934-1935, Cook, A. B., 35-36.
statistics on the productivity of inland waters, Viosca, Percy, Jr., 350-358.
Fish food stocks, warehousing and interviews with meat packers, Fiedler, R. H., and Samson, V. J., 390-395.
supply (bottom and plankton) of Fish Lake, Utah, its abundance and distribution are discussed and tabulated, Hazzard, A. S., 122-128.
Fish foods of Madison lakes as affected by eleven years of chemical treatment, Domogalla, Bernhard, 115-120.

Fish (fresh-water) as affected by crude oil, Wiebe, A. H., 324-330.
grader, new cheap, adjustable type, Needham, P. R., 313-315, 1 plate.
Lake, Utah, preliminary study, Hazzard, A. S., 122-128.
management, creel census as an aid, Eschmeyer, R. W., 222-223.
meal in trout diets, Deuel, Charles R., 161-163.
measurement figured on the basis of how many fish go to the ounce, rather than length of fish, shown to be accurate and handy method, Wales (disc.), 308.
parasites, use of potassium permanganate in control experiments, Wolf, Louis E., 88-100.
policy, preliminary report of committee, Wickliff, E. L., 27-29.
population of Lake Jesse, Nova Scotia, showed a gain in predatory fish over game fish, Smith, M. W., 297-299.
surviving addition of copper sulphate in Lake Jesse, Nova Scotia, Smith, M. W., 109.
(wet basis), spent salmon, canned fish, and fish meals, analysis of protein, fat, carbohydrates and ash content tabulated, 397.
Fisheries Advisory Committee of the Secretary of Commerce, recommendations, Bell, Frank T., 247-248.
investigations in Northeast Brazil, Ihering, Rodolpho von, and Wright, Stillman, 267-271.
Fishery investigation, Federal, and State cooperation, Bell, Frank T., 246-252.
products as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 380-386.
Fishing gear, previous conceptions of selective action of various sizes of mesh in gill and impounding nets exploded by results obtained, Van Oosten, John, 71-73.
in different seasons on Fife Lake, Michigan, analysis of census, Eschmeyer, R. W., 220-222.
in relation to weather conditions, tests showed no correlation between meteorological factors and catch obtained, Eschmeyer, R. W., 217-219.
methods and kinds of bait used during census of Fife Lake, Eschmeyer, R. W., 214-217.
statistics shown to present an entirely wrong picture when "catch per unit time" is employed for nets without regard to the length of time they were left in the water; "The average net lift, corrected for fishing time" should be figured, rather than the "average catch per net per night," Van Oosten, John, 73-75.
Fishway in Mosey River dam, efficient, Cunningham, A. G., 275-276.
Florida, Commission form of conservation government adopted, Denmead, Talbott, 41.
sale of black bass in state prohibited by law, Amsler, Guy, 39.
Food and temperature as affecting the growth of salmon parr in Margaree River, Cape Breton Island, Nova Scotia, Belding, David L., 158.
conservation efficiency in lake, brown, rainbow and brook trout, Tunison, A. V., and McCay, C. M., 366-369.
consumption of fish under natural conditions, how to make daily estimates, Bajkov, A. D., 288-289.
Food for fish, a new mixture used in California hatcheries, Wales, Joseph H., 307-308.
analyses tabulated, 397.
and feeding practices in fish hatcheries, survey, Fiedler, R. H., and Samson, V. J., 376-400.
disastrous results from feeding frozen beef liver in Minnesota hatchery, Surber, Thaddeus (disc.), 402.
dried vs. fresh Wiebe (disc.), 402.
experiments with salmon by-products in feeding young king salmon; mixing coarse flour with spawned-out fish before canning increased the

growth rate and reduced cost of producing the fish when compared with a diet without flour, Donaldson, Lauren R., 165-170.
natural, as studied in fish taken in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 57-69.
natural, preference and selection vs. taking what is available (disc.), 69-70.
natural, 6.05 lbs., wet weight, of Gammarus fasciatus required to produce one pound of brook trout; 6.63 lbs. to produce one pound of rainbow trout, Surber, Eugene W., 300-304.
test of commercial feed combinations, Tunsion, A. V., and McCay, C. M., 369-371.
trout allowed a sufficient amount of commercial food to produce gains comparable with that of other lots tested showed excessive mortality after ten weeks, Deuel, Charles R., 163.
stomach contents of salmon yearling, salmon fry and trout yearling in Gillis' Brook, Scotsville, Cape Breton, analysis tabulated, 153.
Food of trout from lakes in the Klamath River watershed, Doudoroff, Peter, 129-131.
Forage Fish investigations in Michigan, results; experiments in production of red-bellied dace and Menona killifish for pond propagation indicate these fish as suitable additions to the list, Cooper, Gerald P., 132-142.
Foreign Relations Committee, report 1934-1935, Rodd, J. A., 22-24.
Fragilaria, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Fraser River sockeye salmon run, treaty with Canada for its restoration to former size, approved by Canadian Parliament, but not yet ratified by the U. S. Senate, Rodd, J. A., 23.
Freezing plants and cold storage warehouses in the United States and Alaska, tabulated, 391.
Fundulus diaphanus menona, propagation experiments, Cooper, Gerald P., 132-142.
surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Fungus of fish kept under control by spraying method applied to keep algae under control in Madison lakes, Domogalla, Bernhard, 115-120.
Furnace Brook, Vermont's "test stream," statistics of catch, Lord, Russell F., 224-231.
Game-fish, analysis of catch in a Michigan lake, Eschmeyer, R. W., 207-223.
legislative progress chart for 1935, Amsler, Guy, 39.
Gammarus fasciatus, 6.05 lbs. (wet weight) are required to produce one pound of brook trout; 6.63 lbs. to produce one pound of rainbow trout, Surber, Eugene W., 300-304.
limnaeus, abundance in Fish Lake, Utah, Hazzard, A. S., 125.
Gastropoda as food for trout in lakes of the Klamath River watershed, Doudoroff, Peter, 130.
George Washington National Forest, experimental streams established, Davis, H. S., 238-239.
Gerrids surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Gill disease: see **Bacterial** gill.
Gizzard shad: see **Dorosoma** cepedianum.
Glacial acetic acid suggested in large scale treatments for keeping a greater quantity of copper in solution, Surber, Eugene (disc.), 113.
Golden shiner (see also **Notemigonus** crysoleucas), life history observations bearing on their culture, Cooper, Gerald P., 140.
of 23-40 mm. found to feed more on crustaceans than insects; all Crustacea were Cladocera; the insects were all adults and probably taken at the surface; eggs were an important portion of the diet, Ewers, Lela A., and Boesel, M. W., 57-58; tabulated, 62.
Gordon, Seth, elected Secretary-Treasurer American Fisheries Society, 1935, 45-46.
Grader, a new cheap adjustable type, Needham, P. R., 313-315, 1 plate.

Grasshoppers, versus other bait, analysis of catch in Fife Lake, Eschmeyer, R. W., 215-217.
Great Lakes Conservation Council and Lake Erie Advisory Committee, progress in securing uniform regulations reported, Rodd, J. A., 23.
Fisheries, uniform laws and regulations essentially needed, Jackson, Charles E., 26.
Green Mountain National Forest, "test" streams planned, Davis, H. S., 238.
Growth and efficiency of food conversion of lake, brown, rainbow and brook trout, Tunison, A. V., and McCay, C. M., 366-369.
habits of salmon parr, Belding, David L., 157-160.
rate of Cambarus rusticus Girard, influenced by temperature in Ohio fish ponds, Langlois, T. H., 191.
rate of golden shiners, Cooper, Gerald P., 140.
rate, relative of male and female blunt-nosed minnows, Cooper, Gerald P., 137.
Gums, soreness during teething process questioned, Trautman, Milton B., and Hubbs, Carl L., 261-266, 2 figures.
Gyrinids surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Gyrodactylus, treatment of epidemic with potassium permanganate, experimental work, Wolf, Louis E., 88-100.
Habits of Cambarus rusticus Girard in Ohio fish ponds, Langlois, T. H., 189-192.
Halibut, closed season in Northern Pacific fishery ,as instigated by International Fisheries Commission, shown to be beneficial in Report Committee on Foreign Relations, Rodd, J. A., 22-23.
Halifax harbour, transgression of marginal waters over the Scotian Shelf, Leim, A. H., and Hachey, H. B., 279-283.
Hatcheries, survey of food and feeding practices used, Fiedler, R. H., and Samson, V. J., 376-400.
Hatchery disease of Salmonidae, preliminary report on pathology of lesions, symptomatology, etiology and epidemiology, Belding, David L., and Merrill, Beulah, 76-84.
Hatching season of speckled trout in Quebec prolonged from four months to an eight-month period by means of a new method of supplying warm water, Taylor, B. W., 340-341.
Hazzard, A. S., elected Vice-President Division of Aquatic Biology and Physics, 45-46.
Heating water by new, inexpensive method enables winter hatching in Quebec to be extended from four months to eight months, Taylor, B. W., 340-341.
Hectanooga: see **Lake** Hectanooga.
Helioperca incisor specimens taken at four Buckeye Lake stations, stomach contents tabulated, 65.
macrochira, analysis of catch in Fife Lake, Eschmeyer, R. W., 211-212.
Herpobdella punctata surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Herring: see **Sea** herring.
Hewitt's Special Small Trout Food, results of experimental use in New York State Hatchery, Gloversville, Deuel, Charles R., 161-163.
Hirsheimer, Louis C., deceased, 47.
Holopedium found absent in Lake Jesse treated with copper sulphate, and present in control lake, Smith, M. W., 109.
Hormone injection induced extrusion of ripe eggs after 6-10 hours, Ihering, Rodolpho von, and Wright, Stillman, 270.
Horse meat and liver, protein, fat, carbohydrates and ash percentage tabulated, 397.
Human fungous disease spread in Madison lakes, cause discussed, Domogalla, Bernhard, 120-121.

Huntsman, A. G., elected first Vice-President American Fisheries Society, 45-46.
Hyalella knickerbockeri in diet of common sunfish, largemouth bass and yellow perch, in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 59-60.
Hyalotheca, as affected by copper sulphate in Lake Jesse, Smith, M. W., 106-107.
Hyborhynchus notatus, propagation experiments, Cooper, Gerald P., 132-142.
Hydrodictyon control in Madison lakes, Domogalla, Bernhard, 115-120.
Hyland, Carol J., deceased, 47.
Hypophysis extracts injected in Tetragonopterus and Prochilodus induced flow of ripe eggs 6-10 hours later, Ihering, Rodolpho von, and Wright, Stillman, 270.
Ice and cold, research in inland water in winter needed to determine extent of their effect on fish life, Hubbs, Carl L., and Trautman, Milton B., 51-56.
Ictalurus punctatus specimens taken in Buckeye Lake, stomach contents tabulated, 62.
In Memoriam, 47.
Indiana, twenty laws added to its Conservation Code in the 1935 legislature, Amsler, Guy, 38.
Infectious disease of trout described, symptomatology, etiology and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
Insect products as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 388-389.
Insecta in diet of various species of fish in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 57-69.
International Fisheries Commission, regulations for a closed halibut season, beneficial effects shown in report on Foreign Relations Committee, Rodd, J. A., 22-23.
International Pacific Salmon Federation, cooperation prevents duplication of research effort by Canada and the U.S.A., shown in Report Committee on Foreign Relations, Rodd, J. A., 23.
Interstate Sanitation District and Commission, creation by the states of New York, New Jersey and Connecticut authorized by Resolution adopted by Congress (S. J. Res. 159), 32-33.
Intestine of trout as affected by new hatchery disease; pathology, symptomatology and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
Invertebrate forms as affected by copper sulphate in Lake Jesse, Smith, M. W., 109.
Iowa, abandoned mines discharging acid into streams, now sealed, Denmead, Talbott, 31-32.
stream improvement work, Aitken, W. W., 322-323, 2 plates.
Izaak Walton League, new chapters organized in 1934-1935, Denmead, Talbott, 41.
Jesse lake, addition of copper sulphate and results obtained, Smith, M. W., 101-113.
Kansas, one month closed season on black bass adopted, Amsler, Guy, 38.
Kidneys of trout as affected by new hatchery disease; pathology, symptomatology and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
Killifish: see **Fundulus** diaphanes.
Klamath River watershed, food of trout from lakes, Doudoroff, Peter, 129-131.
Labidesthes sicculus specimens taken at three Buckeye Lake stations, stomach contents tabulated, 64.
Lake Champlain and Missisquoi Bay pike-perch fishery as subject of inves-

tigation by International Fact-finding Commission of U. S. A. and Canada, Rodd, J. A., 23.
Dauphin, remarkable productivity, Cunningham, A. G., 275-276.
Edward, Quebec, record of Octomitus salmonis Moore, found in two of the trout examined, Richardson, L. R., 290-292.
Erie Advisory Committee, progress in securing uniform regulations reported, Rodd, J. A., 23.
Erie, western, firmly established and traditionally accepted misconception regarding pollution pointed out, Van Oosten, John, 75.
Fife: see **Fife** Lake.
Fish: see **Fish** Lake.
Hectanooga, as control lake, comparative counts of zooplanktonts with those from Lake Jesse, tabulated, Smith, M. W., 109.
Jesse, addition of copper sulphate and results obtained, Smith, M. W., 101-113.
Jesse, preliminary note on the fish population, Smith, M. W., 297-299.
Mendota, effect of eleven years of chemical treatment on fish and fish foods, Domogalla, Bernhard, 115-120.
Monona, effect of eleven years of chemical treatment on fish and fish foods, Domogalla, Bernhard, 115-120.
Washington, studies on breeding habits of Mylocheilus caurinus, Schultz, Leonard P., 143-147.
Wingra, effect of eleven years of chemical treatment on fish foods, Domogalla, Bernhard, 115-120.
Winnipeg, preservation of whitefish production, Cunningham, A. G., 272-274.
Winnipegosis, history of whitefish depletion during thirty-eight years, Cunningham, A. G., 272-273.
Lake-fed rivers, influence on growth rate of salmon parr, Belding, David L., 159.
Lakes, mountain, in California, set aside to determine the annual production of water of this type, Davis, H. S., 239.
statistics on their productivity, Viosca, Percy, Jr., 350-358.
Langlois, T. H., elected Vice-President Division of Fish Culture, 1935, 45-46.
Largemouth bass: see **Bass,** largemouth.
Laws and regulations concerning changing seasons, limits, etc., in separate states, it is requested that notification thereof be sent the Bureau of Fisheries, Denmead, Talbott, 42.
Leeches (Herpobdella punctata and Macrobdella decora), surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Legislation and Protection, report Vice-President Division of, 1934-1935, 34-70.
Lemna trisulca, distribution and abundance in Fish Lake, Utah, tabulated, 124.
Leonard, J. W., of the Institute for Fisheries Research, Bradt, Glenn W., 254-255.
Lepibema chrysops specimens taken at two Buckeye Lake stations, stomach contents tabulated, 68.
Lesions comparable to those of the eastern bacterial gill disease described by Davis, but with a distinctly different causative organism; histopathological progress of the western type of the disease is discussed, Fish, Frederic F., 85-87.
in form of blebs or blisters in new disease of trout, pathology, symptomatology, etiology, and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
License fees in various states as altered in 1935, Amsler, Guy, 38-39.
Life history of Cambarus rusticus Girard as studied in Ohio fish ponds, Langlois, T. H., 189-192.

Line fishing vs. spearing on Fife Lake, analysis of catch, Eschmeyer, R. W., 219-220.

Live bait, unrestricted use of Tigoma atraria and Cheonda hydrophlox by fishermen proved to be a danger in Fish Lake, Utah, where the chub, thus introduced had become so numerous as to be a distinct annoyance to fly fishermen, Hazzard, A. S., 127.

Liver, addition of 30 per cent to a canned salmon and flour diet proved beneficial in promoting growth without increasing cost of fish produced when compared with a 100 per cent canned salmon and flour diet, Donaldson, Lauren R., 170.

and meal, steam dried vs. air dried, Donaldson, Lauren R., 167.

Liverfluke of sheep, control through extermination of snails by copper sulphate (about two parts per million) caused no loss of food organisms or of algae in the lake, but killed all trout, Hazzard, A. S. (disc.), 114.

Liver of trout as affected by new hatchery disease; pathology, symptomatology and epidemiology discussed, Belding, David L., and Merrill, Beulah, 76-84.

Locke, S. B., elected Vice-President Division of Angling, 45-46.

Log perch: see **Perch,** log.

Lonergan, Senator Augustine, letter addressed to Hon. Harold Ickes, chairman, National Resources Board, quoted in part, 30.

Losses occurring among fish of various sizes after planting, streams in California set aside for this study, Davis, H. S., 239.

Louisiana coast oil well activities offer serious possibilities as polluting agents for oysters, Gowanloch, James Nelson, 293-296.

Macrobdella decora surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.

Macrodon malabaricus in Brazilian waters of uncertain duration, Ihering, Rodolpho von, and Wright, Stillman, 268.

Madison lakes, eleven years of chemical treatment; its effect on fish and fish foods, Domogalla, Bernhard, 115-120.

Management of streams in the National Forests, Davis, H. S., 234-239.

Mansfield bill to prevent pollution of navigable waters not acted on in first Session of 74th Congress, Denmead, Talbott, 31.

Margaree River and tributaries, Cape Breton Island, Nova Scotia, distribution of sucker, salmon and trout, Huntsman, A. G., 152-156.

Maryland passes law giving its commission directionary powers in the matter of changing or regulating seasons, Denmead (disc.), 42.

Meal and liver combinations, steam dried vs. air dried, Donaldson, Lauren R., 170.

fish, cottonseed, meat, salmon egg, in trout diets, Deuel, Charles R., 161-163.

method of preparing air-dried, Donaldson, Lauren R., 167.

Measurement of pollution in fresh water streams, Ellis, M. M., 240-245.

Measuring fish, a new type of grader, cheap and adjustable, Needham, P. R., 313-315, 1 plate.

Meat meal, dried, analysis of protein, fat and ash content tabulated, 397.

meal in trout diets, Deuel, Charles R., 161-163.

packers, interviews with, Fiedler, R. H., and Samson, V. J., 392-395.

products as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 379-380.

Members attending 65th Annual Meeting at Tulsa, Oklahoma, 1935, list of, 15-17.

Membership list, American Fisheries Society: honorary 410; patrons 410-411; life 412; active 413-427; state membership 427-428; clubs, dealers, etc., 428-429; libraries 429; corresponding members 429.

Mendota lake: see **Lake** Mendota.

Menona killifish, life history observations bearing on their culture, Cooper, Gerald P., 141.
Mesocyclops obsoletus, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Mexico, Gulf of, establishment of a scientific laboratory on the Texas coast suggested, Wiebe, A. H., 36.
Michigan, change made in trout season; closed season on white bass abolished, Amsler, Guy, 38.
Fife Lake, analysis of game-fish catch, Eschmeyer, R. W., 207-223.
investigations of forage fish, Cooper, Gerald P., 132-142.
the beaver-trout management program, Bradt, Glenn W., 253-257.
Micrasterias, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Microcystis control in Madison lakes, Domogalla, Bernhard, 115-12-.
controlled with copper sulphate about five parts per million, without any deleterious effect on invertebrate or fish life, Aitken, W. W. (disc.), 113.
Micropterus dolomieu, analysis of catch in Fife Lake, Eschmeyer, R. W., 211-212.
Midge larvae in diet of various species of fish in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 57-69.
Migration, effects of stream obstructions and diversions to be studied in streams set aside for this purpose in California, Davis, H. S., 239.
Milk, dried skim, in trout diets, Deuel, Charles R., 161-163.
Minnow (see also **Blunt-nosed** minnow), propagation experiments, Cooper, Gerald P., 132-142.
stone roller, breeding habits, Smith, Osgood R., 148-151.
versus other bait, analysis of catch in Fife Lake, Eschmeyer, R. W., 215-217.
Monona lake: see **Lake** Monona.
Missisquoi Bay, and Lake Champlain, pike-perch fishery as subject of investigation by International Fact-finding Commission of U. S. A. and Canada, Rodd, J. A., 23.
Morone americana surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Mortality attributable to feeding of fish can be traced to the fact that the product is now always as fresh as it should be, Davis, H. S. (disc.), 400-401.
in hatchery trout caused by new disease, Belding, David L., and Merrill, Beulah, 76-84.
of brook trout excessive when fed sufficient amount of commercial food to produce gains comparable with that of other lots tested, Deuel, Charles R., 163.
Mossy River, Canada, dam and fishway efficient, Cunningham, A. G., 275-276.
Mucus secretion stimulated by chemical bath may cause accumulation of dirt on gills; normally healthy fish without excessive mucous secrtion, not likely to be smothered by excessive silt in water, Hubbs, Carl L. (disc.), 100.
Muskellunge, dentition, Trautman, Milton B., and Hubbs, Carl L., 261-266, 2 figures.
Mylocheilus caurinus, spawning habits, Schultz, Leonard P., 143-147.
Myriophyllum spicatum, distribution, and abundance in Fish Lake, tabulated, 124.
National Forests, stream management, Davis, H. S., 234-239.
National Planning Council of Commercial and Game Fish Commissioners, progress of cooperative program between states and federal government, Jackson, Charles E., 25.

Natural food, extent to which it can be utilized without danger of diminishing returns, experimental streams set aside in California for this study, Davis, H. S., 239.
in diet of artificially propagated fish, methods of obtaining and use in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 389-390.
Nest building habits of Campostoma anomalum Rafinesque, Smith, Osgood R., 148-149.
Net, gill and impounding, previous conceptions of selective action of various sizes of mesh exploded by results obtained, Van Oosten, John, 71-73.
Nevada, districts and seasons reallocated, Amsler, Guy, 38.
New Jersey, creation of Interstate Sanitation District and Commission with the states of New York and Connecticut, authorized by Resolution adopted by Congress (S. J. Res. 159), 32-33.
New York passes law prohibiting sale of muskellunge regardless of where taken and lake trout from all waters of the state except Lakes Erie and Ontario, Amsler, Guy, 39.
creation of Interstate Sanitation District and Commission with the states of new Jersey and Connecticut, authorized by Resolution adopted by Congress (S. J. Res. 159), 32-33.
open season on lake trout shortened, Amsler, Guy, 38.
progress in stream improvement, Greeley, John R., 316-321.
Nitrogen, important factor causing obnoxious water growths in Madison lakes, Domogalla, Bernhard, 115.
was found in almost direct proportion to amount of fertilizer used per week in the pond, Meehean, O. Lloyd, 186.
Nocomis biguttatus, propagation experiments, Cooper, Gerald P., 132-142.
Nomenclature of important North American fishes, resolutions of Committee reported, Hubbs, Carl L., 26-27.
Nominations, report of Committee for election of 1936 officers, 45-46.
North American Council on Fishery Investigations, 1934 meeting on board President Theodor Tissier at Halifax; next meeting to be in Washington, D. C., Sept. 17-19, Rodd, J. A., 24.
North River, Shenandoah Division of the George Washington Forest, to be utilized in an attempt to determine the effect of stream improvement on fishing conditions in general, Davis, H. S., 239.
Northern pike: see **Esox** lucius.
Northern states, tests made on Furnace Brook, Vermont, show that a regulation for a later season and smaller creel-limit for trout-fishing in northern streams would be indicated, Lord, Russell F., 224-231.
Nostoc, distribution and abundance in Fish Lake, Utah, Hazzard, A. S., 124.
Notemigonus crysoleucas auratus, propagation experiments, Cooper, Gerald P., 132-142.
crysoleucas specimens taken at two Buckeye Lake stations, stomach contents tabulated, 62.
crysoleucas surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Notropis cornutus chrysocephalus, propagation experiments, Cooper, Gerald P., 132-142.
Nova Scotia, Lake Jesse, addition of copper sulphate experiments, Smith, M. W., 101-113.
Nutritional requirements of fish (review; composition of fish foods), Fiedler, R. H., and Samson, V. J., 395-397.
of trout, Tunison, A. V., and McCay, C. M., 359-375.
Nutritive value of the blue crab (Callinectes sapidus) and sand crab (Platyonichus ocellatus), Watson, Vernon K., and Fellers, Carl R., 342-349.
Nymphaea advena, new leaves observed after addition of copper sulphate to waters of Lake Jesse, Smith, M. W., 108.

Obstructions and diversions as affecting migration, to be studied in streams set aside for this purpose in California, Davis, H. S., 239.
Octomitus salmonis Moore recorded from Quebec, Richardson, L. R., 290-292.
Odonata as food for trout in lakes of the Klamath River watershed, Northern California, Doudoroff, Peter, 130.
Officers American Fisheries Society for 1934-1935; 1935-1936, 3.
Ohio, abandoned mines discharging acid into streams, now sealed, Denmead, Talbott, 31-32.
fish ponds, notes on the habits of the crayfish, Cambarus rusticus Girard, and its beneficial effects, Langlois, T. H., 189-192.
Oil, crude, effect on fresh-water fish, Wiebe, A. H., 324-330.
pollution in relation to oysters, Gowanloch, James Nelson, 293-296.
Oils, cottonseed, hydrogenated (Crisco) and Salmon oil, utilization by fingerling and yearling brook trout, tabulated, 365.
Oklahoma, law prohibiting sale of black bass now strictly enforced, Denmead, Talbott, 41.
Oncorhynchus keta, air dried salmon meal as food for young king salmon, Donaldson, Lauren R., 165-170.
nerka, experiments show it to be unnecessary for the fish to reach glacial waters to reproduce successfully and an instinctive return to the stream where they were planted, though near a natural spawning stream; a high survival was obtained when unfed fry were planted in a small lake not much above sea level, Dinsmore, A. H., 204-206.
tschawytscha, use of salmon by-products as food in experiments, and results obtained, Donaldson, Lauren R., 165-170.
Oregon, change made in trout season, Amsler, Guy, 38.
Ostracoda in stomachs of fish taken in Buckeye Lake, Ewers, Lela, A. (disc.), 69.
as food for fish: in Mississippi river sloughs ostracods often exceed 20 thousand per square liter of bottom area, yet hardly a trace was found in the food of any of the slough fishes taken, Surber, Eugene (disc.), 70.
Oxygen, dissolved, in measurement of fresh water stream pollution, Ellis, M. M., 243-253.
supply as affected by fertilizer, Wiebe, A. H. (disc.), 36-37.
Oysters, oil pollution affecting, Gowanloch, James Nelson, 293-296.
Pachyurus francisci from Sao Francisco River, Brazil, Ihering, Rodolpho von, and Wright, Stillman, 268.
Pacific Halibut Commission, regulations for a closed season, beneficial effects shown in report on Foreign Relations Committee, Rodd, J. A., 22-23.
salmon research, unprofitable duplication avoided by cooperation between Canada and the U. S. A. through the International Pacific Salmon Federation, Rodd, J. A., 23.
Parasites (see also **Copepod** infection; **Protozoan** diseases), use of potassium permanganate in control work, Wolf, Louis E., 88-100.
Parent stream theory discussed, Belding, David L., 159.
Parker, Winthrop, deceased, 47.
Pelagia noctiluca, hitherto unknown in Canadian waters, prove a transgression of marginal waters over the Scotian Shelf, Leim, A. H., and Hachey, H. B., 279-283.
Pelagic Sealing Treaty of 1911 results in excess of 1,300,000 animals (as per recent U. S. Government estimate) and annual take of well over 50,000 skins of surplus males, Rodd, J. A., 22.
Perca flavescens, analysis of catch in Fife Lake, Michigan, Eschmeyer, R. W., 210-211.
specimens taken at five Buckeye Lake stations, stomach contents tabulated, 67.
surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.

Perch (see also **Perca** flavescens), log, in Buckeye Lake, diet was composed of 67.87 per cent Crustacea, chiefly Cladocera, especially Sida crystallina and Ceriodaphnia reticulata; 30.70 per cent insects, almost altogether midge larvae and pupae, Ewers, Lela A., and Boesel, M. W., 60; tabulated, 67.
white: see **Morone** americana.
yellow (see also **Perca** flavescens), food habits in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 60; tabulated, 67.
Percina caprodes specimens taken at three Buckeye Lake stations, stomach contents tabulated, 67.
Pericardium of trout as affected by new hatchery disease; pathology, symptomatology, epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
Peridinium, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Peritoneum of trout as affected by new hatchery disease; pathology, symptomatology and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
Phillips, J. L., deceased, 47.
Phosphorus, important factor causing obnoxious water growths in Madison Lakes, Domogalla, Bernhard, 115.
Phytoplankton, as affected by copper sulphate in Lake Jesse, Smith, M. W., 106-107.
Pickerel, dentition, Trautman, Milton B., and Hubbs, Carl L., 261-266, 2 figures.
Pike, northern, dentition, Trautman, Milton B., and Hubbs, Carl L., 261-266, 2 figures.
Pike-perch (see also **Stizostedion** vitreum), fishery of Missisquoi Bay and Lake Champlain, subject of investigation by International Fact-finding Commission appointed by U. S. A. and Canada, Rodd, J. A., 23.
Pimelodus clarias from Sao Francisco River, Brazil, Ihering, Rodolpho von, and Wright, Stillman, 268.
Pimephales notatus specimens taken at two Buckeye Lake stations, stomach contents tabulated, 61.
promelas promelas, propagation experiments, Cooper, Gerald P., 132-142.
Pipewort, Eriocaulon articulare, found alive in waters with addition of copper sulphate, Smith, M. W., 108.
Pisgah Division, of Pisgah National Forest, North Carolina, to be utilized for a public demonstration of what can be accomplished by scientific fishery management, Davis, H. S., 239.
Plankton counts of lakes Kegonsa, Mendota, Monona, Waubesa and Wingra, Wisconsin, during eleven years of chemical treatment of Lake Monona, tabulated and plankton counts of Lake Monona, tabulated, Domogalla, Bernhard, 118.
Plant products, analysis of protein, fat, carbohydrates, and ash content tabulated, 397.
as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 387-388.
Platyonichus ocellatus, nutritive value, Watson, Vernon K., and Fellers, Carl R., 342-349.
Pleurataenium, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Plug bait vs. other bait, analysis of catch in Fife Lake, Eschmeyer, R. W., 215-217.
Polluted waters, abnormalities of fish from, possibly caused by high concentrations of carbon dioxide (disc.), Hubbs, Carl L., 203.
"Pollution and Its Consequences,"—a chapter of Senator Hawes' unpublished book—copies available, 32.

Pollution by cities and towns reduced through Public Works Administration, which reports 562 non-federal and 59 federal projects with a total approximate cost of $143,000,000, Denmead, Talbott, 32.
by oil in relation to oysters, Gowanloch, James Nelson, 293-296.
committee report, 1934-1935, Denmead, Talbott, 29-34.
congressional legislation pertaining to, Jackson, Charles E., 25.
in Western Lake Erie, firmly established and traditionally accepted misconception pointed out, Van Oosten, John, 75.
measurement in fresh water streams, Ellis, M. M., 240-245.
oil (disc.), 331.
recommendation that legislation comprise not merely sublethal water purity standards for public waters, but conditions favorable to all stages of the desirable food and game fishes of the particular region and to those aquatic organisms both plant and animal which comprise the food and food chains of these fishes, Denmead, Talbott, 33-34.
resolution concerning, 43-44.
Polyarthra platyptera, as affected by copper sulphate in Lake Jesse, Smith, M. W., 108.
Pomoxis annularis specimens taken at nine Buckeye Lake stations, stomach contents tabulated, 64.
sparoides, analysis of catch in Fife Lake, Eschmeyer, R. W., 212.
Pond culture, underlying requirements and objectives, Mottley, C. McC., 172-178.
fish culture problems at Tulsa municipal fish hatchery, Oklahoma, Aldrich, A. D., 179-183.
plants, Potamogeton americanus, P. filiformis, Ceratophyllum demersum, chemical analysis, tabulated, Meehean, O. Lloyd, 186.
Ponds, dispersal of fertilizing substances, Meehean, O. Lloyd, 184-188.
"Popeye" in new disease of trout, Belding, David L., and Merrill, Beulah, 76-84.
Pork hearts, kidneys, livers, lungs, melts (spleen) analysis of protein, fat, carbohydrates and ash content, tabulated, 397.
Potamogeton americanus, P. filiformis, chemical analyses, Meehean, O. Lloyd, 186.
praelongus, distribution and abundance in Fish Lake, Utah, tabulated, 124.
Potassium permanganate in control of fish parasites, Wolf, Louis E., 88-100
Predatory fish in aquaria, red-bellied dace and blunt-nose minnow added to list, Cooper, Gerald P., 142.
Productivity of inland waters, statistics, Viosca, Percy, Jr., 350-358.
Protein of crab meat, biological value, Watson, Vernon K., and Fellers, Carl R., 345.
Protozoa, Octomitus salmonis Moore recorded from Quebec, Richardson, L. R., 290-292.
Protozoan diseases, Costiasis, identification and treatment, Savage, James, 332-333.
Public Works Administration reports 562 non-federal sewage projects and 59 Federal projects at approximately a total cost of $143,000,000, of which 163 were disposal plants and 458 sewage systems, Denmead, Talbott, 32.
Puget Sound, restoration of sockeye salmon run to its former size through treaty with Canada, approved by Canadian Parliament, but not yet ratified by the U. S. Senate, Rodd, J. A., 23.
Pumpkinseed: see **Eupomotis** gibbosus.
Pungitius pungitius surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Pratt, George D., deceased, 47.
President-elect, speech, Bell, Frank T., U. S. Commissioner of Fisheries, 46.
Presidents, terms of service and places of meeting, 1870-1936, 6-7.

Pribilof Islands, increase in seal herd due to 1911 Pelagic Sealing Treaty with Great Britain, Japan and Russia reported, Rodd, J. A., 22.
Propagation of smallmouth bass in troughs, Kingsbury, O. R., and Royce, Wm. F., 309-312.
Protection and Legislation, report Vice-President Division of, 1934-1935, Amsler, Guy, 37-40.
Quinn, I. T., elected second Vice-President American Fisheries Society, 1935, 45-46.
Racial differences between the parr of different rivers, little evidence left if variations resulting from growth are eliminated, Belding, David L., 159.
Rearing pools in the National Forests, Davis, H. S., 236-237.
Red-bellied dace, life history observations bearing on their culture, Cooper, Gerald P., 141.
Reelfoot Lake, Tennessee, closed season for black bass established, Amsler, Guy, 38.
Regulations, a later season and smaller creel-limit suggested for trout-fishing in northern states, Lord, Russell F., 230.
uniform in Great Lakes area, progress reported, Rodd, J. A., 23.
Reproductive organs of trout as affected by new hatchery disease; pathology, symptomatology and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
Research Committee recommended for American Fisheries Society to see what ought to be done most urgently and assign it to the proper agency, either in the state or the federal government, Wiebe, A. H., 37.
Research work, fishery investigations, Federal and State cooperation, Bell, Frank T., 246-252.
on Pacific salmon, duplication avoided by cooperation of Canada with U. S. A. through the International Pacific Salmon Federation, Rodd, J. A., 23.
pond-rearing as aid in scientific studies, Mottley, C. McC., 175-176.
should include winter field work in inland waters to determine effect of ice and cold on fish life, Hubbs, Carl L., and Trautman, Milton B,, 51-56.
River, size of, as affecting the growth rate of salmon parr, Belding, David L., 158-159.
Rockbass: see **Ambloplites** rupestris.
Run of sockeye salmon, artificially produced, Dinsmore, A. H., 204-206.
Salmincola edwardsii, life history, Savage, James, 335-337, 1 figure.
Salmo fairo, Americanized, yet retain European characteristics, Markus, Henry C., 258-260.
pleuriticus virtually displaced by plantings of rainbow, mackinaw and brook trout in Fish Lake, Utah, Hazzard, A. S., 126.
salar and Salvelinus fontinalis in relation to Catostomus commersonii in Margaree River and tributaries, Cape Breton Island, Nova Scotia, Huntsman, A. G., 152-156.
Salmon as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 383-384.
by-products as food for young king salmon, Donaldson, Lauren R., 165-170.
egg meal, diets containing, gave superior growth at lowest cost in feeding experiments with brook trout at New York State Hatchery, Gloversville, Deuel, Charles R., 162.
king, experimental feeding of young with salmon by-products and results obtained, Donaldson, Lauren R., 165-170.
parr, growth habits, Belding, David L., 157-160.
river system, Nova Scotia; Lake Jesse, experiments in addition of copper sulphate and results obtained, Smith, M. W., 101-113.
sockeye, experiment shows it to be unnecessary for the fish to reach glacial waters to reproduce successfully and an instinctive return to

the stream where they were planted, though a natural spawning stream was near; a high survival was obtained when unfed fry were planted in a small lake not much above sea level, Dinsmore, A. H., 204-206.
sockeye, run to Fraser River and Puget Sound, restoration to its former size through treaty with Canada, approved by Canadian Parliament, but not yet ratified by the U. S. Senate, Rodd, J. A., 23.
Salmonidae, hatchery disease, preliminary report on pathology of lesions, symptomatology, etiology and epidemiology, Belding, David L., and Merrill, Beulah, 76-84.
Salvelinus fontinalis and Salmo salar in relation to Catostomus commersonii in Margaree River and tributaries, Cape Breton Island, Nova Scotia, Huntsman, A. G., 152-156.
fontinalis fontinalis, feeding experiments at the New York State Hatchery at Gloversville, Deuel, Charles R., 161-163.
fontinalis, nutritional studies, Tunison, A. V., and McCay, C. M., 359-375.
fontinalis surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Salyer, J. Clark, of the institute for Fisheries Research, Bradt, Glenn W., 253-254.
Saprolegnia control in Madison lakes, Domogalla, Bernhard, 115-120.
Scales lost while spawning, Smith, Osgood R., 151.
Seapholeberis mucronata, in diet of largemouth bass in Buckeye Lake; most abundant food organism found in stomach contents of brook silverside, Ewers, Lela, A., and Boesel, M. W., 59-60.
Schilbeodes gyrinus specimens taken at two Buckeye Lake stations, stomach contents tabulated, 63.
Schrank, J. J., deceased, 47.
"Schuil Acres," Grand Rapids, Michigan experimental rearing station, results of 1934 propagation experiments, Cooper, Gerald P., 132-138.
Scientific and Common Names of Fishes Committee: see **Committee.**
Scientific and common names of important North American fishes, report, Hubbs, Carl L., 26-27.
Scum control, spraying experiments on Madison lakes, Domogalla, Bernhard, 115-120.
Sea herring as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 382-383.
Seals, annual take of well over 50,000 skins of surplus males reported possible as result of the 1911 Pelagic Sealing Treaty with Great Britain, Japan and Russia, Rodd, J. A., in Report, Committee on Foreign Relations, 22.
Sedges put forth new shoots, but with tips appearing dead, in waters with addition of copper sulphate, Smith, M. W., 108.
Semotilus atromaculatus surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Serrasalmonidae abundant in some Brazilian hydrographic basins, absent in others, Ihering, Rodolpho von, and Wright, Stillman, 268.
Sewage systems and disposal plants installed, according to Public Works Administration reports, 562 projects being non-federal and 59 federal, with a total approximate cost of $143,000,000, Denmead, Talbott, 32.
Shad, gizzard: see **Dorosoma** cepedianum.
Sheep hearts, livers, lungs, analysis of protein, fat, carbohydrates and ash content tabulated, 397.
Shiner (Notemigonus crysoleucas?) analysis of catch in Fife Lake, Michigan, Eschmeyer, R. W., 212.
Shoemaker, Carl D., elected Vice-President Division of Protection and Legislation, 45-46.

Shrimp as food for fish in Federal, State and private hatcheries, Fiedler, R. H., and Samson, V. J., 384-385.
Sida crystallina as food in bluegill and largemouth bass diet in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 59-60.
Silverside: see **Brook** silverside.
Skim milk, dried: see **Milk,** dried skim.
Smolt age of parr, is it uniform? Belding, David L., 160.
Sodium arsenate and copper sulphate vs. crayfish for elimination of vegetation in Ohio fish ponds, Langlois, T. H., 192.
Sodium arsenite, use to inhibit excessive growth of submerged vegetation, Meehean, O. Lloyd, 187.
South Dakota, a 6-inch limit placed on trout and use of live carp as bait prohibited, Amsler, Guy, 38.
Spawning, forced extrusion of eggs induced by injection of hormones, ripe eggs being obtained 6-10 hours after injection, Ihering, Rodolpho von, and Wright, Stillman, 270.
habits of Campostoma anomalum Rafinesque, Smith, Osgood, R., 149-150.
habits of the chub, Mylocheilus caurinus, Schultz, Leonard, P., 143-147.
Spearing vs. line fishing, analysis of catch on Fife Lake, Eschmeyer, R. W., 219-220.
Spears, C. B., deceased, 47.
Spine of trout twisted like the horns of a Rocky Mountain sheep by concentrations of 55-78.5 parts per million of carbon dioxide, Surber, Eugene W., 201; 203.
Spinner bait vs. other bait, analysis of catch in Fife Lake, Eschmeyer, R. W., 215-217.
Spleen of trout as affected by new hatchery disease; pathology, symptomatology and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
Sponges surviving addition of copper sulphate in Lake Jesse, Smith, M. W., 109.
Spraying experiments on Madison lakes during an eleven-year-period, Domogalla, Bernhard, 115-120.
State and Federal cooperation in fishery investigations, Bell, Frank T., 246-252.
Statistics on the productivity of inland waters (master key to better fish culture), Viosca, Percy, Jr., 350-358.
shown to present an entirely wrong picture when "catch per unit time" is employed for nets without regard to the length of time they were left in the water; "the average net lift, corrected for fishing time" should be figured, rather than the "average catch per net per night," Van Oosten, John, 73-75.
Staurastrum, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Steelhead trout: see **Trout,** steelhead.
Stickleback, nine-spines: see **Pungitius** pungitius.
Still-fishing vs. trolling and casting, analysis of catch in Fife Lake, Michigan, Eschmeyer, R. W., 214-217.
Stizostedion vitreum, analysis of catch in Fife Lake, Michigan, Eschmeyer, R. W., 212.
Stocking and artificial propagation in the National Forests, Davis, H. S., 236-237.
of speckled trout from the air, Prevost, Gustave, 277-278.
Stomach analysis of trout from lakes in the Klamath River watershed, North California, tabulated, Doudoroff, Peter, 132.
Stomach contents, method of analyzing, Doudoroff, Peter, 129.
of salmon yearling, salmon fry and trout yearling from Gillis' Brook, Scotsville, Cape Breton, Nova Scotia, tabulated, 153.
Stone roller minnow: see **Minnow,** stone roller.

Stream improvement, management, and surveys by CCC men in the National Forests, Davis, H. S., 234-239.
work in Iowa, Aitken, W. W., 322-323, 2 plates.
Sucker (see also Sweet sucker), in relation to salmon and trout, in Margaree River, and tributaries, Cape Breton Island, Nova Scotia, Huntsman, A. G., 152-156.
Sumner, Francis H., deceased, 47.
Sunfish (see also **Common** sunfish), open season in California extended, Amsler, Guy, 38.
Surirella, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Sweet sucker, food shown to consist of 50 per cent Entomostraca and a more limited quantity of midge larvae and pupae; fish under 37 mm. had eaten no insects; those over 37 mm. had all taken some, Ewers, Lela A., and Boesel, M. W., 57; tabulated, 61.
Symptomatology of new hatchery disease of trout, Belding, David L., and Merrill, Beulah, 76-84.
Tabellaria fenestrata, T. flocculosa, as affected by concentrations of copper sulphate in Lake Jesse, Smith, M. W., 106, 107.
Tadpole catfish, analysis of stomach contents, Ewers, Lela A., and Boesel, M. W., 58; tabulated, 63.
Tankage, dried, analysis of protein, fat and ash content tabulated, 397.
Teeth of the pike, Trautman, Milton B., and Hubbs, Carl L., 261-266, 2 figures.
Temperature and food as affecting the growth of salmon parr in Margaree River, Cape Breton Island, Nova Scotia, Belding, David L., 158.
as affecting the growth rate of Cambarus rusticus in Ohio fish ponds, Langlois, T. H., 191.
glacial waters proved unnecessary for successful reproduction of sockeye salmon, Dinsmore, A. H., 204-206.
Tennessee, closed season on black bass established for Reelfoot Lake, Amsler, Guy, 38.
Texas, establishment of a scientific laboratory on the Gulf Coast for studying coastal waters suggested, Wiebe, A. H., 36.
Thrips, numerous in diet of brook silverside in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 59.
Tigoma atraria and Cheonda hydrophlox introduced into Fish Lake, Utah, by fishermen using them as bait, illustrate the danger of unrestricted use of live bait since the chub has become so numerous as to be a distinct annoyance to fly fishermen, Hazzard, A. S., 127.
Toxic effect of water-soluble fraction of crude oil on fresh-water fish, Wiebe, A. H., 324-330.
Treasurer's report, 1934-1935, Gordon, Seth, 18-20.
Trolling vs. other methods of fishing, analysis of catch in Fife Lake, Eschmeyer, R. W., 214-217.
Troughs, for propagation of smallmouth bass, Kingsbury, O. R., and Royce, Wm. F., 1935, 309-312.
Trout and salmon in relation to the sucker in Margaree River and tributaries, Cape Breton Island, Nova Scotia, Huntsman, A. G., 152-156.
Trout-beaver management program in Michigan, Bradt, Glenn W., 253-257.
Trout, brook (see also **Salvelinus** fontinalis), absorb calcium from the water in which they swim as well as from their diet, Tunison, A. V., and McCay, C. M., 359-375.
experiments show that 6.05 lbs. (wet weight) of Gammarus fasciatus are required to produce a pound of fish, Surber, Eugene W., 300-304.
feeding experiments at the New York State Hatchery at Gloversville, Deuel, Charles R., 161-163.
in Fish Lake, Utah, stomach contents analyzed, Hazzard, A. S., 127.

Trout, brown, even though they are grown to a length of ten or twelve inches in hatcheries, soon gain their natural characteristics when liberated in the stream, Markus, Henry C., 258-260.
Trout, cut throat, eggs as affected by varying concentrations of carbon dioxide, tabulated, 199.
Trout, eggs, effects of carbon dioxide on development, Surber, Eugene W., 194-203.
feeding experiments with natural fish food (Gammarus fasciatus), Surber, Eugene W., 300-304.
feeding in California hatcheries, Wales, Joseph H., 305-308.
fishing in northern states, statistics from Furnace Brook, Vermont's "test stream," show that a regulation for a later season and smaller creel-limit would be advantageous, Lord, Russell F., 224-231.
from lakes in the Klamath River watershed, California, food analyzed, Doudoroff, Peter, 129-131.
killed by copper sulphate (about two parts per million) when used to exterminate snails in lake, though food organisms and algae were not killed, Hazzard, A. S. (disc.), 114.
lake, New York open season shortened, Amsler, Guy, 38.
lesions caused by hatchery disease described; symptomatology, etiology and epidemiology discussed in preliminary report, Belding, David L., and Merrill, Beulah, 76-84.
nutritional requirements, Tunison, A. V., and McCay, C. M., 359-375.
Trout, rainbow, eggs as affected by varying concentrations of carbon dioxide, tabulated, 199.
experiments show that 6.63 lbs. (wet weight) of Gammarus fasciatus are required to produce a pound of fish, Surber, Eugene W., 300-304.
in Fish Lake, Utah, stomach contents analyzed, Hazzard, A. S., 126-127.
Trout season, changes made in Michigan and Oregon, Amsler, Guy, 38.
Trout, speckled, copepod infection, Savage, James, 334-339, 1 figure.
experimental stocking from the air, Prevost, Gustave, 277-278.
hatching and feeding season in Quebec extended from a four to an eight-month period in Quebec, by means of a new cheap method of supplying warm water, Taylor, B. W., 340-341.
Trout, steelhead, life history studies conducted in California streams set aside for this study, Davis, H. S., 239.
Tulsa municipal fish hatchery, Oklahoma, and problems of pond fish culture, Aldrich, A. D., 179-183.
Ulothrix control in Madison lakes, Domogalla, Bernhard, 115-120.
U. S. Bureau of Fisheries actively engaged in pollution investigations, Denmead, Talbott, 32.
recent agreement with Forest Service providing for their close cooperation in putting into effect a scientific program of stream management for the waters of the National Forests, report, Davis, H. S., 234-239.
Varieties, problem of distinct river varieties discussed, Belding, David L., 159.
Vegetation, effect of copper sulphate upon rooted submerged and emergent, Smith, M. W., 107-108.
effectively controlled by introduction of Cambarus rusticus, Langlois, T. H. (disc.), 192.
submerged, chemical analysis showed a fertilizing value of one acre to be equivalent to between 1.35 and two tons of good cottonseed meal, Meehean, O. Lloyd, 187.
Vermont, Furnace Brook, a "test stream," statistics of catch indicate a regulation for later season and smaller creel-limit to be profitable, Lord, Russell F., 224-231.
Walleyed-pike: see **Stizostedion** vitreum.
Washington Lake: see **Lake** Washington.

Washington, state, regulation to provide greater protection of black bass during spawning period, Amsler, Guy, 38.
Water boatmen conspicuous in diet of largemouth bass in Buckeye Lake, Ewers, Lela A., and Boesel, M. W., 60.
hardness, may not be essential to heavy yield of crustacean fish food and trout, Hazzard, A. S., 125.
need for planned utilization, Langlois, T. H., 284-287.
Weather conditions in relation to catch, tests showed no correlation between meteorological factors tested and fish caught, Eschmeyer, R. W., 217-219.
Weed and bathing-beach treatments of Lake Monona, Wisconsin, Domogalla, Bernhard, 119.
West Virginia, abandoned mines discharging acid into streams, now sealed, Denmead, Talbott, 21-32.
Whaling, Multilateral Convention for the Regulation of, signed on behalf of Canada in 1931 at Geneva, ratified at last session of Canadian Parliament, Rodd, J. A., 24.
Wheat shorts and whole wheat flour (graham), analysis of protein, fat, carbohydrates and ash content tabulated, 397.
White bass: see **Bass,** white.
White crappie in Buckeye Lake, diet analyzed, Ewers, Lela A., and Boesel, M. W., 59; tabulated, 64.
Whitefish in Lake Winnipeg, preservation of production, Cunningham, A. G., 272-274.
Willow planting by CCC men in New York State, experiment proving speedy and economical, Greeley, John R., 318-319.
Wingra lake: see **Lake** Wingra.
Winter fish conditions, need for investigation, Hubbs, Carl L., and Trautman, Milton B., 51-56.
hatching in Quebec extended from 4 months to 8 months by use of new inexpensive method of supplying warm water, Taylor, B. W., 340-341.
Wooster, Julian S., deceased, 47.
Worms, versus other bait, analysis of catch in Fife Lake, Eschmeyer, R. W., 215-217.
Xanthidium, as affected by copper sulphate in Lake Jesse, Smith, M. W., 107.
Yellow perch: see **Perch,** yellow.
Zannichellia palustris, distribution and abundance in Fish Lake, Utah, tabulated, 124.
Zooplankton as affected by copper sulphate in Lake Jesse, Smith, M. W., 108-109.

PART II. AUTHOR INDEX

Aitken, W. W.: Iowa stream improvement work, 322-323, 2 plates.

Aldrich, A. D.: The Tulsa municipal fish hatchery and problems of pond fish culture, 179-183.

Amsler, Guy: Report Vice-President Division of Protection and Legislation, 1934-1935, 37-40.

Bajkov, A. D.: How to estimate the daily food consumption of fish under natural conditions, 288-289.

Belding, David L.: Observations on the growth of Atlantic salmon parr, 157-160.

Belding, David L., and **Merrill,** Beulah: Hatchery disease of the Salmonidae, preliminary report, 76-84.

Bell, Frank T., U. S. Commissioner of Fisheries: Inaugural speech as President-Elect of American Fisheries Society, 46.

Federal and State cooperation in fishery investigations, 246-252.

Boesel, M. W., joint author: see **Ewers,** Lela A.

Bradt, Glenn W.: Michigan's beaver-trout management program, 253-257.

Cook, A. B.: Report Division of Fish Culture, 1934-1935, 35-36.

Cooper, Gerald P.: Some results of forage fish investigations in Michigan, 132-142.

Cunningham, A. G.: Preservation of whitefish production in Lake Winnipeg, 272-274.

Remarkable productivity of Lake Dauphin, 275-276.

Davis, H. S.: Mortality attributable to feeding of fish can be traced to the fact that the product is not always as fresh as it should be, 400-401.

Stream management in the National Forests, 1935, 234-239.

Denmead, Talbott: Report, Auditing Committee, 1935, 43.

Preliminary report of the Pollution Study Committee, 1935, 29-34.

Report, Division of Angling, 1935, 40-42.

Deuel, Charles R.: The use of fish meal, cottonseed meal, meat meal, salmon egg meal and dried skim milk in trout diets, 161-163.

Dinsmore, A. H.: Fact vs. theory, 204-206.

Domogalla, Bernhard: Eleven years of chemical treatment of the Madison lakes—its effect on fish and foods, 115-120, map.

Donaldson, Lauren R.: Use of salmon by-products as food for young king salmon, 165-170.

Doudoroff, Peter: Food of trout from lakes in the Klamath River watershed, 129-131.

Ellis, M. M.: Measuring pollution in fresh water streams, 240-245.

Eschmeyer, R. W.: Analysis of the game-fish catch in a Michigan lake, 207-223.

Ewers, Lela A., and **Boesel,** M. W.: The food of some Buckeye lake fishes, 57-69.

Fellers, Carl R., joint author: see **Watson,** Vernon K.

Fiedler, R. H., and **Samson,** V. J.: Survey of fish hatchery foods and feeding practices, 376-400.

Fish, Frederic F.: A western type of bacterial gill disease, 85-87, 4 plates.

Foster, Fred J.: Report, Committee on Nominations, 1935, 45-46.

Gordon, Seth: Report of meeting of the council of American Fisheries Society, Sept. 8, 1935, 21.

Report of the Secretary-Treasurer for 1934-1935, 18-20.

Greeley, John R.: Progress of stream improvement in New York State, 1935, 316-321.

Gowanloch, James Nelson: Pollution by oil in relation to oysters, 293-296.

Hachey, H. B., joint author: see **Leim,** A. H.

Hawes, Senator: "Pollution and its Consequences,"—a chapter in his unpublished book, 32.

Hazzard, A. S.: Preliminary study of exceptionally productive trout water, Fish Lake, Utah, 122-128, map.
Hubbs, Carl L.: Mucus secretion stimulated by chemical bath may cause accumulation of dirt on gills of fish; normally healthy fish without excessive mucous secretion, not likely to be smothered by excessive silt in water, 100.
Report, Committee on the Common and Scientific names of Important North American Fishes, 26-27.
Hubbs, Carl L., and **Trautman,** Milton B.: Need for investigating fish conditions in winter, 51-56.
Hubbs, Carl L., joint author: see **Trautman,** Milton B.
Huntsman, A. G.: The sucker (Catostomus commersonii) in relation to salmon and trout, 152-156.
Ihering, Rodolpho von, and **Wright,** Stillman: Fisheries investigations in Northeast Brazil, 267-271.
Jackson, Charles E.: Report, Committee on Relations with Federal, Provincial and State governments, 1935, 24-26.
Kingsbury, O. R., and **Royce,** Wm. F.: Propagation of smallmouth bass in troughs at South Otselic bass hatchery, 309-312.
Langlois, T. H.: Notes on the habits of Cambarus rusticus in Ohio fish ponds, 189-192.
The need for planned water utilization with aquicultural suggestions, 284-287.
Leim, A. H., and **Hachey,** H. B.: A transgression of marginal waters over the Scotian shelf, 279-283.
Lord, Russell F.: The 1935 trout harvest from Furnace Brook, Vermont's "Test streams," 224-231.
Markus, Henry C.: Americanized brown trout retain European characteristics, 258-260.
McCay, C. M., joint author: see **Tunison,** A. V.
Meehean, O. Lloyd: Dispersal of fertilizing substances in ponds, 184-188.
Merrill, Beulah, joint author: see **Belding,** David L.
Mottley, C. McC.: Objectives in the pond culture of salmonoid game fish, 172-178.
Needham, P. R.: A cheap, adjustable fish grader, 313-315, 1 plate.
Prevost, Gustave: Experimental stocking of speckled trout from the air, 277-278.
Richardson, L. R.: A record of Octomitus salmonis Moore from Quebec, 290-292.
Rodd, J. A.: Report of the Committee on Foreign Relations, 1934-1935, 22-24.
Royce, Wm. F., joint author: see **Kingsbury,** O. R.
Samson, V. J., joint author: see **Fiedler,** R. H.
Savage, James: Copepod infection of speckled trout, 334-339, 1 figure.
Costiasis, identification and treatment, 332-333.
Schultz, Leonard L.: The spawning habits of the chub, Mylocheilus caurinus —a forage fish of some value, 143-147.
Smith, M. W.: A preliminary note on the fish population of Lake Jesse, Nova Scotia, 297-299.
Use of copper sulphate for eradicating the predatory fish population of a lake, 101-113.
Smith, Osgood R.: Breeding habits of the stone roller minnow (Campostoma anomalum Rafinesque), 148-151.
Surber, Eugene W.: Effects of carbon dioxide on the development of trout eggs, 194-203.
Trout feeding experiments with natural food (Gammarus fasciatus), 300-304.
Surber, Thaddeus: Disastrous results from feeding frozen beef liver to trout in Minnesota hatchery (disc.), 402.

Taylor, B. W.: A cheap supply of warm water for winter hatching and feeding of speckled trout, 340-341.
Trautman, Milton B., and Hubbs, Carl L.: When do pike shed their teeth? 261-266, 2 figures.
Trautman, Milton B., joint author: see **Hubbs,** Carl L.
Tunison, A. V.: Economy vs. results obtained in feeding of fish (disc.), 164.
Tunison, A. V., and **McCay,** C. M.: The nutritional requirements of trout, 359-375.
Van Oosten, John: Logically justified deductions concerning the Great Lakes fisheries exploded by scientific research, 71-75.
Viosca, Percy, Jr.: Statistics on the productivity of inland waters: the master key to better fish culture, 350-358.
von Ihering, Rodolpho, and **Wright,** Stillman: Fisheries investigations in Northeast Brazil, 267-271.
Wales, Joseph H.: The feeding of trout in California hatcheries, 305-308.
Watson, Vernon K., and **Fellers,** Carl R.: Nutritive value of the blue crab (Callinectes sapidus) and sand crab (Platyonichus ocellatus Latreille), 342-349.
Wickliff, E. L.: Report of the American Fish Policy Committee, 1935, 27-29.
Wiebe, A. H.: Dried vs. fresh food for fish (disc.), 402.
Report Vice-President Division of Aquatic Biology andd Physics, 1934-1935, 36-37.
The effect of crude oil on fresh-water fish, 324-330.
Type of food eaten due to preference and selection, or to availability? (disc.), 69-70.
Wolf, Louis E.: Use of potassium permanganate in control of fish parasites, 88-100.
Wright, Stillman, joint author: see **Ihering,** Rodolpho von.